

US009636011B2

(12) United States Patent

Sharma et al.

(10) Patent No.: US 9,636,011 B2 (45) Date of Patent: May 2, 2017

(54) SYSTEMS AND METHODS FOR SPECTRALLY DISPERSED ILLUMINATION OPTICAL COHERENCE TOMOGRAPHY

(71) Applicant: Carl Zeiss Meditec, Inc., Dublin, CA (US)

(72) Inventors: Utkarsh Sharma, Dublin, CA (US); Tilman Schmoll, Dublin, CA (US); Xing Wei, Dublin, CA (US)

(73) Assignee: **CARL ZEISS MEDITEC, INC.**, Dublin, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35

U.S.C. 154(b) by 117 days.

(21) Appl. No.: 14/789,889

(22) Filed: Jul. 1, 2015

(65) Prior Publication Data

US 2016/0000320 A1 Jan. 7, 2016

Related U.S. Application Data

- (60) Provisional application No. 62/020,556, filed on Jul. 3, 2014.
- (51) Int. Cl.

 A61B 3/12 (2006.01)

 A61B 3/10 (2006.01)

 G01J 3/45 (2006.01)

 G01J 3/28 (2006.01)
- (58) **Field of Classification Search**CPC A61B 3/102; A61B 2090/3735; A61B
 3/1025; A61B 3/12; A61B 3/1225; A61B
 3/14; A61B 5/0068

USPC 351/206, 246, 205, 200; 356/479, 451, 356/456 See application file for complete search history.

(56) References Cited

U.S. PATENT DOCUMENTS

7,391,520 B2 8,319,974 B2 8,427,654 B2 8,442,284 B2 8,446,593 B1	11/2012 4/2013 5/2013	Zhou et al. Knighton et al. Horn et al. Rogers et al. Ellerbee et al.
8,705,048 B2 2007/0081236 A1	4/2014	Everett et al. Tearney et al.
2007/0001230 711	(Continued)	

OTHER PUBLICATIONS

Blazkiewicz, et al., "Signal-To-Noise Ratio Study of Full-Field Fourier- Domain Optical Coherence Tomography", Applied Optics, vol. 44, No. 36, Dec. 20, 2005, pp. 7722-7729.

(Continued)

Primary Examiner — Hung Dang (74) Attorney, Agent, or Firm — Morrison & Foerster LLP

(57) ABSTRACT

Systems and methods are presented for acquisition and processing of spectrally dispersed illumination optical coherence tomographic data. Light from a source is distributed spectrally on the sample, and each acquisition simultaneously provides partial spectral interference information from multiple locations in the sample. Thus for a given spatial point, a single observation will be of a partial spectrum A-scan. When multiple partial spectrum A-scan observations are made at the same point by shifting the spectrum of light on to the tissue, the point can be observed by the entire broadband spectrum of the light source, thereby making it possible to create a full axial resolution A-scan.

19 Claims, 8 Drawing Sheets

